

Conserved and Divergent Rhythms of Crassulacean Acid Metabolism-Related and Core Clock Gene Expression in the Cactus *Opuntia ficus-indica*^{1[C][W]}

Izaskun Mallona, Marcos Egea-Cortines, and Julia Weiss*

Genetics, Institute of Plant Biotechnology, Technical University of Cartagena, Campus Muralla del Mar, 30202 Cartagena, Murcia, Spain

The cactus *Opuntia ficus-indica* is a constitutive Crassulacean acid metabolism (CAM) species. Current knowledge of CAM metabolism suggests that the enzyme phosphoenolpyruvate carboxylase kinase (PPCK) is circadian regulated at the transcriptional level, whereas phosphoenolpyruvate carboxylase (PEPC), malate dehydrogenase (MDH), NADP-malic enzyme (NADP-ME), and pyruvate phosphate dikinase (PPDK) are posttranslationally controlled. As little transcriptomic data are available from obligate CAM plants, we created an expressed sequence tag database derived from different organs and developmental stages. Sequences were assembled, compared with sequences in the National Center for Biotechnology Information nonredundant database for identification of putative orthologs, and mapped using Kyoto Encyclopedia of Genes and Genomes Orthology and Gene Ontology. We identified genes involved in circadian regulation and CAM metabolism for transcriptomic analysis in plants grown in long days. We identified stable reference genes for quantitative polymerase chain reaction and found that *OfiSAND*, like its counterpart in Arabidopsis (*Arabidopsis thaliana*), and *OfiTUB* are generally appropriate standards for use in the quantification of gene expression in *O. ficus-indica*. Three kinds of expression profiles were found: transcripts of *OfiPPCK* oscillated with a 24-h periodicity; transcripts of the light-active *OfiNADP-ME* and *OfiPPDK* genes adapted to 12-h cycles, while transcript accumulation patterns of *OfiPEPC* and *OfiMDH* were arrhythmic. Expression of the circadian clock gene *OfiTOC1*, similar to Arabidopsis, oscillated with a 24-h periodicity, peaking at night. Expression of *OfiCCA1* and *OfiPRR9*, unlike in Arabidopsis, adapted best to a 12-h rhythm, suggesting that circadian clock gene interactions differ from those of Arabidopsis. Our results indicate that the evolution of CAM metabolism could be the result of modified circadian regulation at both the transcriptional and posttranscriptional levels.

Opuntia species are constitutive Crassulacean acid metabolism (CAM) plants, in which the capacity to induce CAM metabolism is developmentally regulated, with a progression from C₃ to CAM metabolism occurring during cladode development. The CAM cycle is expressed under essentially all growing conditions (Winter et al., 2008). *Opuntia* can take up relatively large amounts of CO₂ with respect to water loss by transpiration (4–10 mmol CO₂ mol⁻¹ water compared with 1–1.5 mmol in C₃ plants), and the annual aboveground dry mass can be increased by 37% to 40% for *Opuntia ficus-indica* when the CO₂ level is doubled (Cui et al., 1993; Nobel et al., 1994).

The CAM type of CO₂ fixation is thought of as integrating into the day/night cycle in four phases. In phase I, nocturnal CO₂ fixation and accumulation of

malic enzyme (ME) occurs in the central vacuole. During phase II, a transition phase early in the light period, malic acid is remobilized from the vacuole and decarboxylated, generating high internal CO₂ concentrations that lead to stomatal closure. Later, in phase III, the released CO₂ is refixed by Rubisco and assimilated via the Calvin cycle of photosynthesis. Finally, in phase IV, the later light period, malic acid stores are exhausted, stomata may open, and CO₂ may be taken up and assimilated directly via Rubisco (Rascher et al., 2001). Key enzymes of CAM metabolism include phosphoenolpyruvate carboxylase (PEPC), which catalyzes CO₂ fixation into phosphoenolpyruvate (PEP) under the formation of oxaloacetate at night; phosphoenolpyruvate carboxylase kinase (PPCK), which controls the phosphorylation state of PEPC; malate dehydrogenase (MDH), required for the reduction of oxaloacetate to malate at night; NADP-ME, catalyzing the decarboxylation of malate during the day in the cytoplasm; and pyruvate phosphate dikinase (PPDK), which is only active in chloroplasts of the malic acid type of CAM species (Cushman and Bohnert, 1997; Scott and Mercer, 1997; Honda et al., 2000) and catalyzes the formation of PEP from pyruvate.

Many photosynthesis-related processes, including starch metabolism, are regulated by a circadian rhythm (Piechulla, 1988, Merida et al., 1999), and the

¹ This work was supported by Fundación Séneca (project ESP-SOL grant to I.M.).

* Corresponding author; e-mail julia.weiss@upct.es.

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors (www.plantphysiol.org) is: Julia Weiss (julia.weiss@upct.es).

[C] Some figures in this article are displayed in color online but in black and white in the print edition.

[W] The online version of this article contains Web-only data.

www.plantphysiol.org/cgi/doi/10.1104/pp.111.179275

circadian clock is related to gas exchange and CAM metabolism through the anticipation of cycles at dawn and dusk (Resco et al., 2009). CAM metabolism displays an endogenous rhythmicity, with circadian oscillations of CO₂ uptake, stomatal conductance, and internal concentrations of CO₂. There is empirical evidence that the circadian transcriptional regulation of *PPCK*, leading to a modulated sensitivity of PEPC to inhibition by malic acid (Nimmo et al., 1987; Hartwell et al., 1999; Taybi et al., 2000), plays a critical role in the control of CAM metabolism (Wyka et al., 2004). *PPCK* is the only CAM-related gene for which a circadian oscillation has been observed at the transcriptional level (Nimmo et al., 1987; Taybi et al., 2000). Phytochrome, acting as a photoreceptor, and the corresponding phytochrome-responsive transcription factors have been suggested to be involved in *PPCK* circadian regulation (Taybi et al., 2000). In general, circadian regulation of enzyme activities during the day/night cycle is mediated largely by posttranslational changes (Cushman et al., 2000) such as the reversible phosphorylation of PEPC by *PPCK* (Chollet et al., 1996; Vidal and Chollet, 1997; Nimmo, 2000), thus connecting PEPC activity to diurnal CAM regulation. *PPDK* activity is light/dark modulated by reversible phosphorylation in both C₃ and C₄ plants (Chastain et al., 2002), and chloroplastic NADP-MDH activity is light regulated via changing NADPH-to-NADP ratios and the thioredoxin redox state (Rebeille and Hatch, 1986; Li et al., 1994). The study of transcriptional patterns of key CAM metabolic enzymes for a plant of the Cactaceae family is novel and offers knowledge about enzyme regulation in CAM plants that can serve as a basis for comparison with other CAM types and C₄ plants.

In *Arabidopsis* (*Arabidopsis thaliana*), a central circadian oscillator complex is formed by the genes *CIRCADIAN CLOCK ASSOCIATED1/LATE ELONGATED HYPOCOTYL* (*CCA1/LHY*) acting together with *TIMING OF CAB EXPRESSION1* (*TOC1*) in a feedback-regulating system, operating throughout the life of the plant in a diversity of tissues (Locke et al., 2005, 2006). It controls stomatal conductance rhythms, hypocotyl elongation, and flowering time via the integration of photoperiodic timing and temperature (Somers et al., 1998; Strayer et al., 2000; Ding et al., 2007). Products of these genes interact with those from a set of other clock genes, like *PSEUDO-RESPONSE REGULATOR9* (*PRR9*), a critical component of the circadian system and an integrator of temperature signals (Salomé and McClung, 2005). Although the circadian clock has been extensively studied in *Arabidopsis* (de Montaigne et al., 2010), the genetic contributors and their expression patterns have not been described in more than a few species (Beales et al., 2007; Iwamoto et al., 2009; Hayes et al., 2010). Studying these circadian oscillator genes in CAM plants should help us to understand the circadian regulation of key CAM metabolic enzymes.

As little transcriptomic data are available from obligate CAM plants, we developed a large collection of *Opuntia* ESTs. It is accessible through the *Opuntia*-

ESTdb, providing information about clusters, annotations, tandem repeats, and metabolic pathways. As a direct application, we used the *Opuntia*ESTdb to retrieve the sequences of CAM-related and circadian clock genes. There are multiple isoforms for photosynthetic genes with differing functions in plants with C₃, C₄, and CAM metabolism, and as expected, we obtained several contigs for each gene studied. We isolated a set of possible reference genes as a prerequisite for validating transcriptomic changes by quantitative (Q) real-time PCR. Reference genes, traditionally thought of as housekeeping genes, are needed in order to normalize mRNA levels between different samples, correcting for differences in the amount of starting material or RNA isolation efficiencies, thus allowing for exact interpretation of gene expression data (Vandesompele et al., 2002). Housekeeping genes, whose products are involved in basic functions needed for cell maintenance, were initially assumed to be constitutively expressed, but the expression levels of these genes can vary among tissues or cells (Mallona et al., 2010) and may be modified by environmental conditions. It can therefore be concluded that no gene can be presumed a priori to be "constitutively expressed" (Czechowski et al., 2005; Gutierrez et al., 2008). The selection of appropriate reference genes for the types of tissues and circumstances under investigation is required for accurate gene expression studies. We identified *OfiSAND* and *OfiTUB* as suitable genes for the normalization of transcription levels in a variety of tissues. After validating the adequacy of reference genes, we performed Q-PCR relative quantification and found several canonical gene expression profiles, including rhythmic expression of *OfiPPCK* and *OfiTOC1* and arrhythmic expression of *OfiPEPC* and *OfiMDH*. Surprisingly *OfiNADP-ME*, *OfiPPDK*, *OfiCCA1*, and *OfiPRR9* adopted a 12-h cycle with two peaks during the day, indicating significant differences in the structure of chronobiological regulation in CAM plants.


RESULTS

EST Annotation Overview

The sequencing of cDNA derived from RNA pools of various tissues (Fig. 1) generated a total of 604,176 ESTs with an average read length of 344 bp (Supplemental Table S2). Clustering of these ESTs produced a total of 43,066 contigs and 407,253 unassembled singletons with average lengths of 612 and 1,385 bp, respectively. Annotation against the National Center for Biotechnology Information (NCBI) nonredundant database showed that 29,835 contigs produced the lowest BLAST hit scores against sequences from a CAM species (69.3%) as listed by Sayed (2001), whereas 1,015 were closer to those of *Arabidopsis* (2.4%).

The *Opuntia*ESTdb Web interface includes a searchable database through BLASTN, TBLASTN, BLASTX,

Figure 1. Organs and developmental stages used for RNA extraction prior to EST sequencing and real-time Q-PCR. A, Three-month-old seedling. B, Young cladode. C, Fully grown cladode. D, Young flower bud. E and F, Flower buds at anthesis. G and H, Fruits. Bars = 1 cm, except for C, where bar = 5 cm.



TBLASTX, and BLASTP assembled contigs and singletons. Preexistent ESTs from CAM species were recovered from The Institute for Genomic Research assemblies and included in the database to allow for more comprehensive searches. Contig sequence recovery includes functional annotation fetching of the annotations obtained from its RefSeq/UniProtKB putative orthologs. Thus, Kyoto Encyclopedia of Genes and Genomes (KEGG) Orthologs and Pathways, PubMed publications, ExPaSy, InterPro, and Gene Ontology (GO) annotations, Tandem Repeats, and UniProtKB cellular locations and keywords are retrieved if present. KEGG Pathways including *O. ficus-indica* contigs are offered as highlighted maps. Additional information on data analysis and functional categorization can be found in the Supplemental Data, including a flow chart of database construction and its applications (Supplemental Fig. S1), the comparative distribution of a selection of functional categories between the classified genes from the Arabidopsis genome and the Opuntia EST clusters as detected by WEGO (Supplemental Fig. S2A), the distribution of functional categories among contigs showing the 10 most common GO terms in the EST database for each of the three ontology domains, molecular function, cellular component, and biological process (Supplemental Fig. S2B), and the KEGG pathway for carbon fixation in photosynthetic organisms with highlighted orthologs from the Opuntia ESTdb (Supplemental Fig. S3).

Validation of Reference Genes for Q-PCR in *O. ficus-indica*

The cDNA samples obtained from six different organ types of *O. ficus-indica* were used for real-time Q-PCR with the eight primer pairs designed for amplification of the candidate reference genes (Supplemental Table S1). For each run, the cycle threshold (CT) value (the number of cycles required for normalized fluorescence to reach a threshold), reaction efficiency, and product melting temperature were evaluated. Single melting

peaks for all reference gene products indicated single-product amplification.

The CT values associated with each technical replicate were quality checked, and those differing from the average by more than one cycle were discarded. Values for 90% of the technical replicates differed by less than one cycle. Figure 2 as well as Supplemental Table S3 show the normalization results. Global results from stability and coefficient of variation values of qBasePlus and stability values from geNorm, as well as a pairwise variation cutoff of 0.15, indicate that for all single organs, a combination of two reference genes is sufficient and necessary for normalizing target gene expression (Fig. 2B). Combining two organ types requires two (for cladodes sampled at both day and night) and three (for ovules and fruits) reference genes, while for combining all tissue types, no pairwise variation value underneath the cutoff limit could be determined. Figure 2A indicates the reference gene ranking for single and mixed organs based on cutoff values of 0.5 for stability and 0.2 for coefficient of variation. Supplemental Figure S4 shows the relative quantities of reference gene expression for each biological replicate.

Identification of CAM and Circadian Clock Homologs

Several genes encoding CAM-related enzymes (e.g. *PEPC*, *ME*, and *MDH*) belong to small gene families (Cushman and Bohnert, 1989a, 1989b; Cushman et al., 1993; Honda et al., 2000). For example, up to four members of *PEPC* exist in *Mesembryanthemum crystallinum* and *Kalanchoe blossfeldiana*, with one or two CAM-specific isoforms (Cushman and Bohnert, 1997). As the Arabidopsis genome is thoroughly annotated, we used Arabidopsis ab initio to estimate the gene family structure of all genes analyzed. We found several contigs that fulfilled the criteria of homology and performed phylogenetic analysis of the contigs against available sequences from C_3 , C_4 , and CAM plants (Fig. 3).

Several isoforms of *PPCK* have been associated with C_4 , CAM, or housekeeping C_3 types. There are two




Figure 2. Reference gene expression analysis. Organ types are as follows: YC, young cladodes; CD, cladodes sampled during the day; CD, cladodes sampled at night; OV, ovule tissue of fully developed flowers before anthesis; FR, ripe fruits. A, Reference gene expression stability heat map. geNorm stability (M) and qBasePlus stability and coefficient of variation (CV) expression values are shown for single and mixed organs. Each heat map spot represents a reference gene-organ or organ mix combination; spots with values below a stability threshold are represented in green (meaning reference gene suitability), whereas those above that threshold are plotted in magenta (not suitable). Specific thresholds are 0.2 for qBasePlus coefficient of variation values and 0.5 for geNorm and qBasePlus stability values. A superposed heat map, combining results from the three previous layers, is labeled as Global. Green reflects combinations that had three out of three suitable stability measures, yellow reflects two out of three, orange reflects one out of three, and magenta reflects none. B, Minimum number of genes necessary for reliable and accurate normalization. GeNorm pairwise variation values are computed by an algorithm that measures pairwise variation ($V_n/n + 1$) between two sequential normalization factors, NFn and $NFn + 1$, where n is the number of genes involved in the normalization factor. A pairwise variation threshold of 0.15 is plotted. [See online article for color version of this figure.]

PPCK genes in *Arabidopsis*; using *PPCK1* as query, we identified three contigs and two singletons. Phylogenetic analysis of the *O. ficus-indica* sequences revealed two singletons and two contigs clustered together in a branch that includes genes from CAM and non-CAM plants, whereas a fifth clone clusters with *Arabidopsis* and other plants, including C_4 plants (Fig. 3A). The data from this phylogenetic reconstruction did not show any obvious separation of CAM and non-CAM sequences (i.e. those from the C_3 species *Beta vulgaris*, *Solanum lycopersicum*, and the CAM species *Kalanchoe daigremontiana* and *M. crystallinum*). A different *O. ficus-indica* sequence grouped together with *Arabidopsis*, *Zea mays*, *Theobroma cacao*, and *Glycine max*, reinforcing the scattered pattern of *PPCK* sequences.

We found 10 contigs of *PEPC* using *AtPEPC2*, a member of the small gene family comprising four loci in *Arabidopsis*, as query. The tree topology shows four clusters of *OfiPEPC* genes, including one close to *AtPEPC4*, and another group highly homologous to the CAM plant *Ananas comosus* (Fig. 3B). There is yet another group that clusters together with CAM- and C_3 -type clones from *M. crystallinum*, suggesting again a complex evolutionary history for the *Opuntia* gene family that cannot be resolved purely by sequence similarity criteria.

A total of 17 loci code for MDH in *Arabidopsis*, with apoplasmic, chloroplasmic, mitochondrial, and peroxisomal enzyme activities. The sequence used as query encodes an enzyme that is apoplasmic as well as chlo-

roplasmic, with NAD^+ or $NADP^+$ (chloroplasmic) as acceptor. It has been suggested that combined activities of cytosolic NAD^+ -MDH and chloroplasmic $NADP$ -MDH may be involved in nocturnal malic acid formation in CAM plants (Cushman, 1993). We identified five *OfiMDH* contigs and four *OfiMDH* singletons (Fig. 3C). We observed two major branches, with all *O. ficus-indica* clones clustering within one branch that also includes MDH from the CAM species *Welwitschia mirabilis*. The second main branch includes MDH isoforms of C_4 plants like *Z. mays* and *Sorghum vulgare* as well as one from the CAM species *M. crystallinum* and C_3 plants. The clustering of all *O. ficus-indica* clones indicates high similarity among possible isoforms in *Opuntia*. Nevertheless, CAM-specific isoforms fail to cluster together.

Malate decarboxylation in Cactaceae species, which belong to the ME group of CAM plants, mainly occurs via the activity of cytoplasmic $NADP$ -ME (Scott and Mercer, 1997), although substantial activities of mitochondrial NAD -ME might also contribute to decarboxylation in CAM plants (Cushman and Bohnert, 1997). Four *NADP*-ME isoforms exist in *Arabidopsis*, cytosolic except for *NADP-ME4*, which is chloroplasmic. We used as query the cytosolic *Arabidopsis* *NADP-ME2* and identified 10 *O. ficus-indica* contigs satisfying the homology cutoff score (Fig. 3D). These contigs cluster on two branches. One branch includes five *O. ficus-indica* contigs as well as the CAM isoform of *NADP-ME* from




Figure 3. Phylogenetic analyses of PPCK (A), PEPC (B), MDH (C), NADP-ME (D), and PPDK (E). Bootstrapping values (percentage from 10,000 sampled trees) are shown at the branch points. *O. ficus-indica* clones labeled with "contig" or "F" plus letter/number code are contigs and singletons, respectively, identified from the EST database. Clones labeled as *O. ficus-indica* contig used were chosen for expression analysis. [See online article for color version of this figure.]

M. crystallinum. The second branch including *O. ficus-indica* clones clusters with isoforms of C_4 (*Z. mays* and *Flaveria trinervia*) and CAM (*Aloe arborescens*) plants. A third major branch contains isoforms from both C_3 and C_4 plants. Tree topology indicates that NADP-ME isoforms from CAM plants tend not to gather with those from C_3 plants but are more similar to C_4 isoforms.

The Arabidopsis genome contains one chloroplastic *PPDK* gene, while in C_4 plants, both C_4 and non- C_4 isoforms have been identified (Chastain et al., 2002). Six *O. ficus-indica* contigs and three singletons are highly homologous to *PPDK* from Arabidopsis, clustering on two branches (Fig. 3E). One branch also contains *PPDK* of *M. crystallinum* as well as isoforms of C_4 and C_3 *Flaveria* species. The second major cluster carries *O. ficus-indica* sequences as well as *PPDK* isoforms from several C_4 plants (*Z. mays*, *Saccharum officinarum*, and *Sorghum bicolor*) as well as *Oryza sativa* and *Triticum aestivum*. The *AtPPDK* was the most divergent sequence in the *PPDK* tree.

In summary, except for *NADP-ME*, which separates *O. ficus-indica* clones clearly from those of C_3 species, the CAM-related genes do not clearly cluster separately from those encoding C_3 or C_4 isoforms or their host species.

There are five *PRR* genes in the Arabidopsis genome that form three clades, corresponding to *AtPRR1-TOC1*, *PRR3* and *PRR7*, and *PRR5* and *PRR9* (Takata et al., 2010). Our phylogenetic reconstruction (Fig. 4A) shows two branches, one containing the *O. ficus-indica* genes clearly isolated from the rest and a second containing Arabidopsis, *G. max*, *Populus trichocarpa*, and two *O. ficus-indica* genes, suggesting rapid evolution of a group of *PRR* genes in the genus *Opuntia*. An additional group of genes involved in circadian regulation are those encoding MYB transcription factors, out of which *CCA1* and *LHY* are considered the morning oscillators (de Montaigu et al., 2010). We found only three genes with high homology to Arabidopsis *LHY* and *CCA1* (Fig. 4B), all three clustering closer to *CCA1* than to *LHY*. In contrast, the *CCA1* gene from *M. crystallinum* was more similar to that of *Vitis vinifera*, indicating a lack of CAM specificity in the clustering of these genes.

As our results showed no clustering of CAM, C_3 , or C_4 genes, we used the homolog closest to the Arabidopsis query sequence for further analysis.

Temporal Gene Expression Analysis by Q-PCR

We followed the expression pattern of three genes coding for enzymes active in the cytoplasm during the night, *OfiPEPC*, *OfiPPCK*, and *OfiMDH*, and two-light active enzymes, *OfiNADP-ME* and *OfiPPDK* (Fig. 5). *OfiPEPC* and *OfiMDH* expression patterns did not fit to a circadian rhythm (Supplemental Table S3). In contrast, *OfiNADP-ME* and *OfiPPDK* showed significant adjustment to a 12-h rhythm, while *OfiPPCK* oscillated with a 24-h periodicity. *OfiPPCK* expression was highest about 3 h after the beginning of the dark period; *OfiNADP-ME* and *OfiPPDK* showed increases in expression during the morning and again in the afternoon approaching the




Figure 4. Phylogenetic analysis of TOC (A) and PRR (B) genes. Bootstrapping and clone labeling are as described for Figure 3. [See online article for color version of this figure.]

dark period. For core clock genes, *OfiTOC1* adapted to a circadian rhythm with lowest expression during the morning and the acrophase at dusk. In contrast, homologs to the Arabidopsis clock genes *OfiCCA1* and *OfiPRR9* both followed statistically significant 12-h oscillations. Peak expression of these two genes coincided in the morning and again at dusk. This 12-h period has not been observed in Arabidopsis, where *CCA1* and *PRR9* follow a circadian rhythm. Similar to *TOC* expression in Arabidopsis, *OfiTOC1* showed an evening peak slightly later than the evening peaks of *CCA1* and *PRR9*.

DISCUSSION

EST Cluster Generation and Sequence Annotation

ESTs provide a shortcut to the transcribed portions of the genome, allowing a rapid screen for novel genes based on BLAST searches. The sequencing of *O. ficus-*




Figure 5. Gene expression chronograms of CAM and circadian rhythm pathways during the day/night cycle in cladodes of *O. ficus-indica*. Three normalized, relative gene expression quantities corresponding to three biological replicates were collected every 4 h (circles). Oscillation (represented as a continuous curve) was modeled, via Fourier analysis, decomposing it in two harmonic or sinusoidal components: one circadian (24-h harmonic) and one ultradian (12 h). Dotted curves indicate harmonic adjustment. The solid line indicates the proposed model including all significant harmonics. a.u., Arbitrary units. [See online article for color version of this figure.]

indica cDNA performed here generated 604,176 sequences. Sixty-seven percent of these sequences were unassembled singletons, not clustering with any other sequences. These stand-alone ESTs may be from regions of low coverage; nevertheless, similar numbers were reported for other EST collections (Cervigni et al., 2008). The *OpuntiaESTdb* presented here is, to our knowledge, the first online database of a plant species performing CAM metabolism whose sequences can be freely accessed and downloaded, with valuable information about whole-plant gene expression.

From the contigs whose sequences showed homology to those present in the public databases, 29,835 (69%) had significant matches with categorized proteins, and putative functions were assigned on this basis. Nonmatching contigs might indicate specific roles for the products of these sequences in constitutive CAM plants. The *Arabidopsis* genome is predicted to contain more than 25,000 protein-coding genes and the *Z. mays* genome at least 32,540 (Schnable et al., 2009). Assuming a number in *O. ficus-indica* similar to that in *Z. mays*, the identified 29,835 protein-coding

genes represents close to 90% of the *O. ficus-indica* gene complement (Segura et al., 2007).

Validation of Reference Genes for Q-PCR in *O. ficus-indica*

The identification of *O. ficus-indica* gene homologs to well-categorized protein-coding genes from other organisms opens up the possibility for broad-scale transcriptional analysis. Quantitative analysis relies upon reference genes with essentially invariant transcript levels in different tissues under different conditions; the *OpuntiaESTdb* allowed us to identify *O. ficus-indica* gene homologs to known reference genes. Because CO_2 absorption changes from C_3 to CAM metabolism during cladode development (Winter et al., 2008), basing our analysis on both young and fully grown cladodes allowed us to analyze CO_2 uptake-related gene expression during this developmental step. We also used adult cladodes sampled during day and night as prerequisite for analyzing the transcript levels of genes showing circadian oscillation. Finally, the determination

of optimal reference genes in floral and fruit tissue is crucial for investigating the genes involved in flower and fruit development, for example those related to parthenocarpic fruit development (Weiss et al., 1993) and fruit ripening (Collazo-Siqués et al., 2003). Our results show that the use of two reference genes is sufficient and necessary for single and mixed organs except for the combination “ovules plus fruits,” which requires three reference genes. Interestingly, the *SAND* protein (*SAND*), proposed as a reference gene for Arabidopsis (Czechowski et al., 2005), proved the most reliable reference gene for the set of organs examined here, followed by *b-Tubulin6* (*TUB*).

Gene Identification and Phylogenetic Analysis

Phylogenetic analysis of CAM-related and circadian clock-related homologs from *O. ficus-indica* showed no specific clustering with genes and/or CAM-specific isoforms of other CAM plants. With the exception of *O. ficus-indica* clones for NADP-ME, which did not seem to cluster with C₃ isoforms of this enzyme, all the genes under investigation had contigs and singletons in mixed branches that included C₃ and C₄ isoforms.

Irrespective of their photosynthetic metabolism, C₄ and CAM plants have C₃ housekeeping as well as C₄- and CAM-specific isoforms of photosynthetic enzymes. It has been postulated that C₄ photosynthesis arose in a multistep process starting with the duplication of genes for ubiquitous metabolic enzymes, which then acquired specific functions in a light-, development-, cell-, and tissue-specific manner (Ku et al., 1996; Besnard et al., 2003; Monson, 2003). Our phylogenetic analysis for PEPC showed that, in the case of the CAM plants *M. crystallinum* and *K. blossfeldiana*, the C₃ and CAM isoforms were more similar within the species than they were to C₃ and CAM isoforms of the other CAM species. We conclude that *O. ficus-indica* CAM-specific isoforms cannot be identified based on phylogenetic analysis. Lack of clear grouping among CAM-type isoforms, as observed in our analysis, has also been reported by Taybi et al. (2000) for PPCK from *M. crystallinum*, whose most closely related PPCKs were those from *S. lycopersicum* followed by *Kalanchoe fedtschenkoi* and Arabidopsis. Concerning PPDK, Chastain et al. (2011) found only minor differences between C₃ and C₄ isoforms from Arabidopsis and *Z. mays* in key catalytic and regulatory properties. Those authors propose a transition from a functionally C₃ isoform into a C₄ pathway enzyme involving only minor changes in enzyme properties. The suggested mechanism could account for the close relationships between CAM and C₃ and C₄ isoforms observed in our phylogenetic analysis. CAM metabolism is widespread, occurring within 33 plant families, and water stress is proposed as the ultimate selective factor for terrestrial CAM plants (Keeley and Rundel, 2003). The widespread systematic occurrence hints at multiple independent

evolutionary events (Smith and Winter, 1996) and might explain the phylogenetic tree structures observed here.

Rhythmic Expression of CAM and Clock Genes

We studied the temporal expression pattern of *O. ficus-indica* genes homologous to circadian clock- and CAM-related genes using the identified reference genes. The objectives were (1) to compare the expression patterns of several CAM-related genes in the cactus *O. ficus-indica* with those known from other CAM species, with special focus on the circadian oscillation pattern, and (2) to determine whether central oscillator components in the circadian cycle known from Arabidopsis show the typical circadian rhythm of transcription in *O. ficus-indica*.

While the enzymatic machinery required to perform CAM is present in all higher plants, activities of the key enzymes are much higher in CAM and C₄ plants, and it is proposed that they have acquired diurnal patterns of expression and regulation to meet nighttime CO₂ fixation and daytime decarboxylation of C₄ acids (Cushman and Bohnert, 1997). It has also been suggested that posttranslational control mediates circadian regulation of enzyme activities during the day/night cycle, while transcriptional, posttranscriptional, and translational control is primarily responsible for the buildup of CAM-related enzymes (Cushman et al., 2000).

In the case of PEPC, short-term responses are post-translationally controlled, as shown for *M. crystallinum* (Li and Chollet, 1994) and *K. fedtschenkoi* (Carter et al., 1996) as well as in *Crassula argentea*, where two PEPC configurations exist and modification of the configuration involves phosphorylation and dephosphorylation (Lambers, 2008). This observation is in accordance with our data showing that the *OfiPEPC* homolog did not show major transcriptional changes over a 24-h period (Fig. 5; Supplemental Tables S4 and S5).

The only CAM-related enzyme for which a circadian oscillation rhythm in transcription has been reported in different CAM species is PPCK. A diurnal oscillation in transcript abundance was reported for the constitutive CAM plant *K. fedtschenkoi* (Hartwell et al., 1999) and the facultative CAM plant *M. crystallinum* (Taybi et al., 2000). The circadian transcription of the gene for PPCK, which leads to a modulated sensitivity of PEPC to inhibition by malic acid, is proposed to be the key clock system responsible for circadian regulation of CAM metabolism (Nimmo et al., 1987; Hartwell et al., 1999; Wyka et al., 2004). Our studies confirm the diurnal expression pattern of PPCK in *O. ficus-indica*, although the acrophase was observed at dusk, while highest transcript levels in *M. crystallinum* and *K. fedtschenkoi* occur toward dawn. Thus, activation of PEPC by PPCK seems to be regulated at the transcriptional level and might represent a crucial control point in CAM metabolism in *O. ficus-indica*, but with significant differences from the system reported for *M. crystallinum* and *K. fedtschenkoi*.

The enzyme MDH catalyzes the conversion of oxaloacetate to malate. In *M. crystallinum*, a gene encoding NADP⁺-dependent MDH was isolated that shows a salt stress-induced expression pattern, indicating its participation in the CO₂ fixation pathway during CAM (Cushman, 1993). Circadian regulation of MDH activity was reported to be posttranslational, as chloroplastic NADP-MDH activity is light regulated in response to the NADPH/NADP⁺ ratio and the thioredoxin redox state (Rebeille and Hatch, 1986; Li et al., 1994). The selected *OfiMDH* gene, similar to *OfiPEPC*, did not follow a circadian expression rhythm but rather a constitutive transcription pattern during a 24-h course, suggesting a reliance on posttranscriptional regulation.

Species of the Cactaceae family belong to the ME type of CAM plant, and malate decarboxylation mainly occurs via the activity of cytoplasmic NADP-ME (Scott and Mercer, 1997). NADP-ME plants require PPDK in the chloroplast for gluconeogenic recovery of PEP. Its activity is under dark/light regulation by reversible phosphorylation in *Z. mays* (Budde et al., 1985; Chastain et al., 2002), involving posttranslational regulation through the PPDK regulatory protein (Chastain et al., 2002, 2011). It was reported for *M. crystallinum* that transcript levels of NADP-ME increase 8- to 10-fold in response to salt stress in the leaves as CAM metabolism is induced (Cushman, 1992). In our studies, transcription of the two enzymes adjusted to a 12-h rhythm, and acrophase almost coincided for these enzymes. A rise in transcript levels starts at the beginning of CAM phase III, when CO₂ is released from malate, refixed by Rubisco, and assimilated via the Calvin cycle of photosynthesis, and the resulting pyruvate is converted into PEP. A second rise in expression during phase IV might be related to an increased demand of PEP for CO₂ fixation. Our results show that in *O. ficus-indica*, transcription of genes for PPDK and NADP-ME changes during the day and these changes, together with posttranslational activations, might be relevant for the adaptation of enzyme activity during the CAM cycle.

Control of gene expression by the circadian clock involves some central oscillator genes. *CCA1/LHY* together with *TOC1* form a core oscillator complex in Arabidopsis. *CCA1* and *LHY* bind to the *TOC1* promoter and negatively regulate its expression, while *TOC1* positively regulates *CCA1/LHY* together with other genes, including *GIGANTEA*, *EARLY FLOWER-ING3 (ELF3)*, *ELF4*, and *LUX ARRITHMO* (McClung, 2006). *PRR9* acts as an oscillator necessary for the clock to respond to temperature signals, anticipating diurnal cold stress and initiating a stress response by mediating cyclic expression of stress response genes, including *DREB1/CBF* (Salomé and McClung, 2005; Nakamichi et al., 2009). *PRR9* is positively regulated by *CCA1/LHY*, but it negatively regulates *CCA1/LHY* (McClung, 2006).

Our results show that of the *O. ficus-indica* homologs to Arabidopsis central oscillator genes *CCA1*, *TOC1*, and *PRR9*, only *OfiTOC1* shows circadian rhythmicity. *OfiTOC1*, similar to its expression profile in Arabidop-

sis, has its acrophase at the onset of the dark period. In Arabidopsis, the morning-activated oscillator *CCA1* shows an opposite rhythm, as *CCA1* inhibits *TOC1* while *TOC1* induces *CCA1*, creating an autoregulatory feedback loop (Locke et al., 2006). In *O. ficus-indica*, *OfiCCA1* and *OfiPRR9* seem to adapt to an ultradian 12-h rhythm and not to the 24-h cycle seen in Arabidopsis, with peak expression during the morning and a second expression peak at dusk before the rise in *OfiTOC1* expression. Our results indicate that homologs to key oscillator genes from Arabidopsis do not build the same interconnected autoregulatory loops in *O. ficus-indica*, pointing to the involvement of other key oscillator genes in this CAM species. It remains to be determined if the evolution of CAM occurred by changes in the setup of the circadian clock rather than by a modification of downstream genes in their cis-regulatory regions.

MATERIALS AND METHODS

cDNA Library Construction and Sequencing

A normalized cDNA library was constructed from RNA isolated from six individuals of the species *Opuntia ficus-indica*, collected from different sites in the Mediterranean southeast of Murcia, Campo de Cartagena, Spain, where these plants are typically planted by commercial growers as small populations alongside other crops such as citrus or *Vitis vinifera*. Total RNA was extracted from seedlings, young cladodes, fully grown cladodes sampled during the day and night, young flower buds, fully developed flowers, and ripe fruits (Fig. 1). Total RNA was purified using the Nucleospin RNA Plant Kit (Macherey-Nagel), which includes a recombinant DNase I treatment in order to eliminate genomic DNA contamination. Sequencing was performed by Eurofins MWG Operon applying 454 sequencing technology.

EST Sequence Analysis

Raw sequence data were clustered using CAP3 with a stringency level of 95% similarity per 100 bp; clipping of poor regions regarded base quality values and similarity information (Huang and Madan, 1999). Assembled EST data sets were compared using BLASTX (e value < 1e-10) with the NCBI nonredundant database in order to identify putative homologs. In the case of significant BLASTX, the hit with minimal e value was ascribed to the contig description, and thus a RefSeq and NCBI GI identifier was assigned to each contig. These accessions were mapped to KEGG Orthology and GO. KEGG Orthology terms were assigned to KEGG Genes based on cross-database links to GI identifiers and compared with Arabidopsis (*Arabidopsis thaliana*) as background using KOBAS (for KEGG Orthology-Based Annotation System; Wu et al., 2006). Relevant metabolic reactions were plotted with the KEGG Pathway exploring tool. RefSeq accessions were mapped to UniProtKB identifiers and thereafter to GO using DAVID (for Database for Annotation, Visualization, and Integrated Discovery; Dennis et al., 2003; Da Wei Huang and Lempicki, 2008). GO abundance was compared with the Arabidopsis background and represented with WEGO (Ye et al., 2006). Orthologs for the selected pathways were obtained from the collection of plant annotated sequences available at the GO consortium, and they were accessed through the European Bioinformatics Institute QuickGO interface (Binns et al., 2009). Protein sequences involved in those processes were retrieved from the QuickGO GOA files and compared via TBLASTN against the *Opuntia*ESTdb (e value cutoff of 1e-10). For each significant *Opuntia* sequence, only the BLAST record hit with the lowest e value was taken, thus avoiding multiple assignments.

The database architecture runs on Ubuntu Linux with an apache2 HTTP server and MySQL as the database management system. Pages can be requested through several Common Gateway Interface python scripts. The source code is available at the server's page.

Identification of Genes with Stable Patterns of Gene Expression

We selected a list of eight genes considered to exhibit stable patterns of expression and therefore suitable for use as internal controls for Q-PCR (Andersen et al., 2004; Czechowski et al., 2005; Nicot et al., 2005; Mallona et al., 2010). The list consisted of *Actin-11* (*ACT*), *Cyclophilin-2*, *Elongation factor 1a* (*EF1*), *Ubiquitin GTP-binding protein RAN1*, *SAND*, *Ribosomal protein S13*, and *TUB* (Supplemental Table S1). Highly homologous contigs or singletons to the corresponding Arabidopsis identifiers were retrieved from the *Opuntia*ESTdb using TBLASTN. Primers were designed using the application PRIMEGENS version 2 (which discards cross-hybridizations within the EST database; Srivastava et al., 2008), resulting in an amplicon size between 103 and 154 bp (Supplemental Table S1).

Total RNA was extracted as described above using young cladodes (6–8 cm length), fully grown cladodes sampled during the day and at night, ovule tissue of fully developed flowers before anthesis, and ripe fruits. Tissue samples were taken from one of the populations used for RNA sequencing. Three independent samples, representing pools of three samples per plant, were taken for each tissue and time course. RNA concentration and purity were estimated from the ratio of absorbance readings at 260 and 280 nm, and RNA integrity was tested by gel electrophoresis. cDNA synthesis was performed from equal amounts of total RNA using Moloney murine leukemia virus reverse transcriptase (SuperScript III First-Strand Synthesis SuperMix; Invitrogen). Real-time PCR was carried out in an Mx3000P Q-PCR system (Stratagene) using a SYBR Green-based PCR assay (with 5-carboxy-X-rhodamine as the optional reference dye; Brilliant II SYBR Green Q-PCR Master Mix; Stratagene), applying a PCR protocol of 95°C for 5 min, 40 cycles of 95°C for 30 s, 60°C for 30 min, and 72°C for 30 s, followed by a standard dissociation protocol to assess product uniqueness. All assays were performed in three technical replicates for each of the three biological replicates and included nontemplate controls.

For data mining and statistical analysis, PCR amplification kinetics were analyzed to calculate crossing points and reaction efficiencies with the qpcR R package version 1.3-3 (Spiess et al., 2008). The efficiency value (E) was defined as $E = F_n / (F_n - 1)$, in which n is determined as the 20% value of the normalized fluorescence at the maximum of the second derivative curve. Data generated for potential internal controls were quality tested and checked for expression stability using qBasePlus version 1.5 (Hellemans et al., 2007), which includes stability and coefficient of variation determination as well as the classical $geNorm$ parameters such as stepwise stability and pairwise variation (Vandesompele et al., 2002).

Identification of *O. ficus-indica* Homologs and Phylogenetic Analysis

For each candidate circadian-regulated gene, we searched the *Opuntia*ESTdb via TBLASTN (e value $< 1e-10$) using an Arabidopsis accession as the query sequence. In order to analyze sequence dissimilarities between the putative *O. ficus-indica* homologous sequences and those corresponding to other C_4 , C_3 , and CAM species, we employed the same Arabidopsis sequence with TBLASTN against the NCBI nonredundant database. Significant hits were recovered, and further accessions based on literature searches were included in the phylogenetic work flow.

All sequences were translated and aligned using a progressive alignment algorithm with default parameters (Feng and Doolittle, 1987) and clustered via neighbor joining. Internal node support was assessed using 10,000 bootstrap replicates. Sequence alignment and clustering were performed with the CLC bioinformatic suite version 6. The accessions employed were as follows.

(1) Query: Arabidopsis PPCK1 AT1G08650; C_3 plants: Arabidopsis CDPK AT1G08650, *Glycine max* PPCK3 AY144184.1, *G. max* PPCK2 AY144182.1, *G. max* PPCK AY143660.2, *Theobroma cacao* EST EST01262, *Solanum lycopersicum* PPCK AF203481.1, *Beta vulgaris* PPCK AJ309171.1, *Solanum tuberosum* PPCK AF453448, *Oryza sativa* PPCK1 AK101080.1, *O. sativa* PPCK2 XM46683.1, and *O. sativa* PPCK3 AK066885; CAM plants: *Kalanchoe fedtschenkoi* PPCK AF162661, *Mesembryanthemum crystallinum* PPCK AF158091, *Kalanchoe pinnata* PPCK EF157816.1, and *Kalanchoe daigremontiana* PPCK EF157817.1; C_4 plants: *Zea mays* PPCK4 AY911416, *Z. mays* PPCK1 (C_4 type) AY911413, *Z. mays* PPCK3 AY911413, *Z. mays* PPCK2 (non- C_4 type) AY911415, and *Flaveria bidentis* PPCK AB065100.1.

(2) Query: Arabidopsis PEPC2 AT2G42600; C_3 plants: *Arachnea hypogea* PEPC EV391629.1, *G. max* PEPC4 AY563044.1, *Sesuvium portulacastrum* PEPC

DQ655705.1, *S. lycopersicum* PEPC1 AJ243416.1, *Medicago sativa* PEPC M83086.1, *Citrus sinensis* PEPC EF058158.2, Arabidopsis PPC4 AT1G68750, Arabidopsis PPC1 AT1G53310, *Gossypium hirsutum* PEPC1 AF008939.1, *Ricinus communis* PEPC1 EF634319.1, Arabidopsis PPC3 AT3G14940, and *Flaveria pringlei* PEPC1 Q01647; CAM plants: *Opuntia streptacantha* PEPC EX720539, *M. crystallinum* PPC1 (CAM type) X13660.1, *M. crystallinum* PPC2 P16097.1, *Kalanchoe blossfeldiana* PPC1 (CAM type) CAA61083.1, *K. blossfeldiana* PPC2 (CAM type) CAA61084.1, *K. blossfeldiana* PPC3 (C_3 type) CAA61085.1, and *K. blossfeldiana* PPC4 (C_3 type) CAA61086.1; C_4 plants: *Flaveria trinervia* PPC-A (C_4 type) P30694, *F. trinervia* PEPC1 (non- C_4 type) Q9FV66, *Ananas comosus* PEPC DV190685, *A. comosus* similar to PEPC DT338073, *Z. mays* PEPC1 NP 001105418.1, *Z. mays* PEPC2 (C_3 type) X61489.1, *Z. mays* PEPC2 (C_4 type) X03613.1, *Sorghum vulgare* PEPC (C_3 type) X59925.1, and *S. vulgare* PEPC (C_4 type) X17379.1.

(3) Query: Arabidopsis MDH AT3G47520.1; C_3 plants: *R. communis* MDH XM002514704.1, *Brassica rapa* MDH FJ208590, *Citrus junos* MDH DQ901430.1, *F. pringlei* MDH P36444, and *Nicotiana tabacum* MDH AJ006574.1; CAM plants: *Welwitschia mirabilis* EST DT584016 and *M. crystallinum* MDH chloroplast Q05145; C_4 plants: *Z. mays* MDH cytoplasm Q08062, *Z. mays* MDH (C_4 type) P15719, *S. vulgare* MDH (C_4 type) P17606, *S. vulgare* MDH Q43830, *F. trinervia* MDH (C_4 type) P22178, and *F. trinervia* MDH Q42737.

(4) Query: Arabidopsis cytosolic NADP-ME2 NM121205.3; C_3 plants: *N. tabacum* cytosolic NADP-ME DQ923118.2, *F. pringlei* NADP-ME AF288921.1, Arabidopsis cytosolic NADP-ME3 AT5G25880, *O. sativa* NADP-ME AY435404.1, *Hordeum vulgare* NADP-ME EU977175.1, Arabidopsis cytosolic NADP-ME1 AT2G19900, *R. communis* ME XM002514184, *S. lycopersicum* chloroplast NADP-ME AF001269.1, and Arabidopsis NADP-ME4 chloroplast AAF68118.1; CAM plants: *M. crystallinum* NADP-ME (C_4 type) CAA45772.1, *Aloe arborescens* NADP-ME (C_3 type) AB005808.1, and *A. arborescens* NADP-ME (C_4 type) BAA24950.1; C_4 plants: *Z. mays* NADP-ME (non- C_4 type) U39958.1, *Z. mays* NADP-ME (C_4 type) J05130.1, *F. trinervia* NADP-ME (non- C_4 type) EU118291.1, and *F. trinervia* NADP-ME (C_4 type) X61304.1.

(5) Query: Arabidopsis PPDK AT4G15530; C_3 plants: *O. sativa* PPDK NM 001062042, *Triticum aestivum* EST CK208491.1, and *F. pringlei* PPDK Q42736; CAM plant: *M. crystallinum* PPDK CAA55143; C_4 plants: *Saccharum officinarum* PPDK AF194026.1, *Sorghum bicolor* PPDK AY268138.1, *Z. mays* PPDK1-1 (C_4 type) P11155-1, *Z. mays* PPDK1-2 (non- C_4 type) P11155-2, *F. bidentis* PPDK (C_4 type) Q39735, *F. trinervia* PPDK (C_4 type) P22221-1, and *F. trinervia* PPDK (non- C_4 type) P22221-4.

The clock genes used for phylogenetic reconstruction were as follows.

(1) Query: Arabidopsis TOC1 AT5G61380 and Arabidopsis APRR9 AT2G46790; C_3 plants: *Pisum sativum* PRR59 FJ609179.1, *Castanea sativa* TOC1 AY611028.1, *G. max* TOC1 EU076435.1, *Populus trichocarpa* PRR9 XM002330094.1, Arabidopsis AIP1 AJ251086.1, Arabidopsis TOC1 AT5G61380, Arabidopsis APRR1 AB041530.1, *B. rapa* PRR1a GU219472.1, Arabidopsis TOC1-like AF272040.1, *B. rapa* PRR9 GU219476.1, Arabidopsis APRR3 NM125403.2, Arabidopsis APRR5 NM122355.3, and *P. trichocarpa* PRR XM002318466.1; CAM plant: *M. crystallinum* TOC1 AY371288.1; C_4 plants: *Z. mays* TOC1 HQ003892.1 and *Z. mays* PRR1 HM452303.1.

(2) Query: Arabidopsis CCA1 AT2G4630; C_3 plants: *Populus nigra* LHY1 AB429410.1, *R. communis* EST XM002515047.1, Arabidopsis MYP-TF AT2G46830, *T. aestivum* LHY HQ222606.1, *G. max* LCL2 EU076434.1, *C. sativa* LHY AY611029.1, Arabidopsis LHY NM_179237.1, *S. lycopersicum* EST BT012912.1, and *Vitis vinifera* EST XM002267684.1; CAM plant: *M. crystallinum* CCA1 AY371287.1; C_4 plants: *Z. mays* LHY NM001138057 and *Z. mays* CCA1 HM452304.1.

Temporal Gene Expression Analysis by Q-PCR

Contigs highly homologous to Arabidopsis genes involved in CAM metabolism and the circadian cycle were retrieved from the *Opuntia*ESTdb using TBLASTN (Supplemental Table S1). Primers for an amplicon length between 266 and 349 bp were designed using the application PRIMEGENS version 2 in order to ensure unique hybridization within the EST database.

As plant material, 1-year-old *O. ficus-indica* cladodes, clonally derived from one mother plant, were grown in the greenhouse in a substrate of perlite: coconut fiber:floral substrate (ratio of 1:1:1) and watered as required. Cladodes were transferred to Sanyo MRL350 growth chambers for adaptation for 1 week and programmed for a light regime of 16 h of light and 8 h of darkness at day/night temperatures of 22°C/15°C and a photosynthetically active photon flux density of 250 $\mu\text{E s}^{-1} \text{m}^{-2}$. Transversal, circular samples of 1 cm diameter were taken every 4 h from three plants. Total RNA extraction and cDNA synthesis as well as real-time PCR were performed as described above using the same PCR cycling conditions.

CT and efficiency values for each single PCR run were used as input for the qBasePlus software, which computed relative quantities against a normalization factor based on *EF1* and *ACT*, which satisfied the geNorm stability measure. In order to detect rhythmic patterns, relative quantities were processed with the El Temps tool (<http://www.el-temps.com/>), which included a Fourier analysis comprising one or two harmonics. By this means, periodicity, Euler's coefficients (a, b), acrophase, amplitude, phase, significance, Snedecor's F, power, and cumulated power were obtained. Details about Fourier and Cosinor analyses are listed in Supplemental Tables S4 and S5.

The OpuntiaESTdb is freely available at <http://srvgen.upct.es/opuntia/index.html> (user: opuntia; password: Opuntia ficus-indica). All questions, comments, and requests may be sent by e-mail to julia.weiss@upct.es.

Supplemental Data

The following materials are available in the online version of this article.

Supplemental Figure S1. Work flow scheme of OpuntiaESTdb.

Supplemental Figure S2. EST functional categories.

Supplemental Figure S3. KEGG pathways for "carbon fixation in photosynthetic organisms."

Supplemental Figure S4. Reference gene expression profiling.

Supplemental Table S1. Primer information and amplicon characteristics.

Supplemental Table S2. Characteristics of the OpuntiaESTdb sequences.

Supplemental Table S3. Optimal genes for expression quantification in individual and mixed organs.

Supplemental Table S4. Fourier analysis.

Supplemental Table S5. Cosinor analysis.

ACKNOWLEDGMENTS

We thank Luis Pedro García and SEDIC for aid in conducting the computation-intensive part of this work as well as the Web server setup. Marta Pawluczyk is acknowledged for comments on the manuscript. Special thanks to Judith Strommer for help with editing.

Received April 28, 2011; accepted June 14, 2011; published June 15, 2011.

LITERATURE CITED

- Andersen CL, Jensen JL, Ørntoft TF (2004) Normalization of real-time quantitative reverse transcription-PCR data: a model-based variance estimation approach to identify genes suited for normalization, applied to bladder and colon cancer data sets. *Cancer Res* **64**: 5245–5250
- Beales J, Turner A, Griffiths S, Snape JW, Laurie DA (2007) A pseudo-response regulator is misexpressed in the photoperiod insensitive Ppd-D1a mutant of wheat (*Triticum aestivum* L.). *Theor Appl Genet* **115**: 721–733
- Besnard G, Pinçon G, D'Hont A, Hoarau JY, Cadet F, Offmann B (2003) Characterisation of the phosphoenolpyruvate carboxylase gene family in sugarcane (*Saccharum* spp.). *Theor Appl Genet* **107**: 470–478
- Binns D, Dimmer E, Huntley R, Barrell D, O'Donovan C, Apweiler R (2009) QuickGO: a Web-based tool for Gene Ontology searching. *Bioinformatics* **25**: 3045–3046
- Budde RJ, Holbrook GP, Chollet R (1985) Studies on the dark/light regulation of maize leaf pyruvate, orthophosphate dikinase by reversible phosphorylation. *Arch Biochem Biophys* **242**: 283–290
- Carter PJ, Fewson CA, Nimmo GA, Nimmo HG, Wilkins MB (1996) Role of circadian rhythms, light and temperature in the regulation of phosphoenolpyruvate carboxylase in Crassulacean acid metabolism. In K Winter, JAC Smith, eds, *Crassulacean Acid Metabolism: Biochemistry, Ecophysiology and Evolution*. Springer, Berlin, pp 46–52
- Cervigni GD, Paniego N, Díaz M, Selva JP, Zappacosta D, Zanazzi D, Landerreche I, Martelotto L, Felitti S, Pessino S, et al (2008) Expressed sequence tag analysis and development of gene associated markers in a near-isogenic plant system of *Eragrostis curvula*. *Plant Mol Biol* **67**: 1–10
- Chastain CJ, Failing CJ, Manandhar L, Zimmerman MA, Lakner MM, Nguyen TH (2011) Functional evolution of C4 pyruvate, orthophosphate dikinase. *J Exp Bot* **62**: 3083–3091
- Chastain CJ, Fries JP, Vogel JA, Randklev CL, Vossen AP, Dittmer SK, Watkins EE, Fiedler LJ, Wacker SA, Meinhover KC, et al (2002) Pyruvate, orthophosphate dikinase in leaves and chloroplasts of C₃ plants undergoes light-/dark-induced reversible phosphorylation. *Plant Physiol* **128**: 1368–1378
- Chollet R, Vidal J, O'Leary MH (1996) Phosphoenolpyruvate carboxylase: a ubiquitous, highly regulated enzyme in plants. *Annu Rev Plant Physiol Plant Mol Biol* **47**: 273–298
- Collazo-Siqués PV, Valverde ME, Paredes-López O, Guevara-Lara F (2003) Expression of ripening-related genes in prickly pear (*Opuntia* sp.) fruits. *Plant Foods Hum Nutr* **58**: 317–326
- Cui M, Miller PM, Nobel PS (1993) CO₂ exchange and growth of the Crassulacean acid metabolism plant *Opuntia ficus-indica* under elevated CO₂ in open-top chambers. *Plant Physiol* **103**: 519–524
- Cushman JC (1992) Characterization and expression of a NADP-malic enzyme cDNA induced by salt stress from the facultative Crassulacean acid metabolism plant, *Mesembryanthemum crystallinum*. *Eur J Biochem* **208**: 259–266
- Cushman JC (1993) Molecular cloning and expression of chloroplast NADP-malate dehydrogenase during Crassulacean acid metabolism induction by salt stress. *Photosynth Res* **35**: 15–27
- Cushman JC, Bohnert HJ (1989a) Nucleotide sequence of the gene encoding a CAM specific isoform of phosphoenolpyruvate carboxylase from *Mesembryanthemum crystallinum*. *Nucleic Acids Res* **17**: 6745–6746
- Cushman JC, Bohnert HJ (1989b) Nucleotide sequence of the Ppc2 gene encoding a housekeeping isoform of phosphoenolpyruvate carboxylase from *Mesembryanthemum crystallinum*. *Nucleic Acids Res* **17**: 6743–6744
- Cushman JC, Bohnert HJ (1997) Molecular genetics of Crassulacean acid metabolism. *Plant Physiol* **113**: 667–676
- Cushman JC, Meiners MS, Bohnert HJ (1993) Expression of a phosphoenolpyruvate carboxylase promoter from *Mesembryanthemum crystallinum* is not salt-inducible in mature transgenic tobacco. *Plant Mol Biol* **21**: 561–566
- Cushman CJ, Taybi T, Bohnert HJ (2000) *Advances in Photosynthesis*, Vol 9. Kluwer Academic Publishers, Dordrecht, The Netherlands
- Czechowski T, Stitt M, Altmann T, Udvardi MK, Scheible WR (2005) Genome-wide identification and testing of superior reference genes for transcript normalization in *Arabidopsis*. *Plant Physiol* **139**: 5–17
- Da Wei Huang B, Lempicki R (2008) Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc* **4**: 44–57
- de Montaigu A, Tóth R, Coupland G (2010) Plant development goes like clockwork. *Trends Genet* **26**: 296–306
- Dennis G Jr, Sherman BT, Hosack DA, Yang J, Gao W, Lane HC, Lempicki RA (2003) DAVID: database for annotation, visualization, and integrated discovery. *Genome Biol* **4**: 3
- Ding Z, Doyle MR, Amasino RM, Davis SJ (2007) A complex genetic interaction between *Arabidopsis thaliana* TOC1 and CCA1/LHY in driving the circadian clock and in output regulation. *Genetics* **176**: 1501–1510
- Feng DE, Doolittle RF (1987) Progressive sequence alignment as a prerequisite to correct phylogenetic trees. *J Mol Evol* **25**: 351–360
- Gutierrez L, Mauriat M, Guénin S, Pelloux J, Lefebvre JF, Louvet R, Rusterucci C, Moritz T, Guérineau F, Bellini C, et al (2008) The lack of a systematic validation of reference genes: a serious pitfall undervalued in reverse transcription-polymerase chain reaction (RT-PCR) analysis in plants. *Plant Biotechnol J* **6**: 609–618
- Hartwell J, Gill A, Nimmo GA, Wilkins MB, Jenkins GI, Nimmo HG (1999) Phosphoenolpyruvate carboxylase kinase is a novel protein kinase regulated at the level of expression. *Plant J* **20**: 333–342
- Hayes KR, Beatty M, Meng X, Simmons CR, Habben JE, Danilevskaya ON (2010) Maize global transcriptomics reveals pervasive leaf diurnal rhythms but rhythms in developing ears are largely limited to the core oscillator. *PLoS ONE* **5**: e12887
- Hellemans J, Mortier G, De Paepe A, Speleman F, Vandesompele J (2007) qBase relative quantification framework and software for management and automated analysis of real-time quantitative PCR data. *Genome Biol* **8**: R19
- Honda H, Akagi H, Shimada H (2000) An isozyme of the NADP-malic enzyme of a CAM plant, *Aloe arborescens*, with variation on conservative amino acid residues. *Gene* **243**: 85–92
- Huang X, Madan A (1999) CAP3: a DNA sequence assembly program. *Genome Res* **9**: 868–877

- Iwamoto M, Higo K, Takano M (2009) Circadian clock- and phytochrome-regulated Dof-like gene, Rdd1, is associated with grain size in rice. *Plant Cell Environ* 32: 592–603
- Keeley JE, Rundel PW (2003) Evolution of CAM and C4 carbon-concentrating mechanism. *Int J Plant Sci* 164: S55–S77
- Ku MS, Kano-Murakami Y, Matsuoka M (1996) Evolution and expression of C4 photosynthesis genes. *Plant Physiol* 111: 949–957
- Lambers H (2008) *Plant Physiological Ecology*. Springer, New York
- Li B, Chollet R (1994) Salt induction and the partial purification/characterization of phosphoenolpyruvate carboxylase protein-serine kinase from an inducible Crassulacean-acid-metabolism (CAM) plant, *Mesembryanthemum crystallinum* L. *Arch Biochem Biophys* 314: 247–254
- Li D, Stevens FJ, Schiffer M, Anderson LE (1994) Mechanism of light modulation: identification of potential redox-sensitive cysteines distal to catalytic site in light-activated chloroplast enzymes. *Biophys J* 67: 29–35
- Locke JCW, Kozma-Bognár L, Gould PD, Fehér B, Kevei E, Nagy F, Turner MS, Hall A, Millar AJ (2006) Experimental validation of a predicted feedback loop in the multi-oscillator clock of *Arabidopsis thaliana*. *Mol Syst Biol* 2: 59
- Locke JCW, Southern MM, Kozma-Bognár L, Hibberd V, Brown PE, Turner MS, Millar AJ (2005) Extension of a genetic network model by iterative experimentation and mathematical analysis. *Mol Syst Biol* 1: 2005.0013
- Mallona I, Lischewsky S, Weiss J, Hause B, Egea-Cortines M (2010) Validation of endogenous genes as controls for quantitative real-time PCR during leaf and flower development in *Petunia hybrida*. *BMC Plant Biol* 10: 4
- McClung CR (2006) Plant circadian rhythms. *Plant Cell* 18: 792–803
- Merida A, Rodriguez-Galan JM, Vincent C, Romero JM (1999) Expression of the granule-bound starch synthase I (Waxy) gene from snapdragon is developmentally and circadian clock regulated. *Plant Physiol* 120: 401–410
- Monson RK (2003) Gene duplication, neofunctionalization, and the evolution of C4 photosynthesis. *Int J Plant Sci* 164: 43–54
- Nakamichi N, Kusano M, Fukushima A, Kita M, Ito S, Yamashino T, Saito K, Sakakibara H, Mizuno T (2009) Transcript profiling of an *Arabidopsis* PSEUDO RESPONSE REGULATOR arrhythmic triple mutant reveals a role for the circadian clock in cold stress response. *Plant Cell Physiol* 50: 447–462
- Nicot N, Hausman J-E, Hoffmann L, Evers D (2005) Housekeeping gene selection for real-time RT-PCR normalization in potato during biotic and abiotic stress. *J Exp Bot* 56: 2907–2914
- Nimmo GA, Wilkins MB, Fewson CA, Nimmo HG (1987) Persistent circadian rhythms in the phosphorylation state of phosphoenolpyruvate carboxylase from *Bryophyllum fedtschenkoi* leaves and its sensitivity to inhibition by malate. *Planta* 170: 408–415
- Nimmo HG (2000) The regulation of phosphoenolpyruvate carboxylase in CAM plants. *Trends Plant Sci* 5: 75–80
- Nobel PS, Andrade JL, Wang N, North GB (1994) Water potentials for developing cladodes and fruits of a succulent plant, including xylem-versus-phloem implications for water movement. *J Exp Bot* 45: 1801–1807
- Piechulla B (1988) Plastid and nuclear messenger-RNA fluctuations in tomato leaves: diurnal and circadian rhythms during extended dark and light periods. *Plant Mol Biol* 11: 345–353
- Rascher U, Hütt MT, Siebke K, Osmond B, Beck F, Lüttge U (2001) Spatiotemporal variation of metabolism in a plant circadian rhythm: the biological clock as an assembly of coupled individual oscillators. *Proc Natl Acad Sci USA* 98: 11801–11805
- Rebeille F, Hatch MD (1986) Regulation of NADP-malate dehydrogenase in C4 plants: relationship among enzyme activity, NADPH to NADP ratios, and thioredoxin redox states in intact maize mesophyll chloroplasts. *Arch Biochem Biophys* 249: 171–179
- Resco V, Hartwell J, Hall A (2009) Ecological implications of plants' ability to tell the time. *Ecol Lett* 12: 583–592
- Salomé PA, McClung CR (2005) PSEUDO-RESPONSE REGULATOR 7 and 9 are partially redundant genes essential for the temperature responsiveness of the *Arabidopsis* circadian clock. *Plant Cell* 17: 791–803
- Sayed O (2001) Crassulacean acid metabolism 1975–2000, a check list. *Photosynthetica* 39: 339–352
- Schnable PS, Ware D, Fulton RS, Wei F, Pasternak S, Liang C, Zhang J, Fulton L, Graves TA, et al (2009) The B73 maize genome: complexity, diversity, and dynamics. *Science* 326: 1112–1115
- Scott T, Mercer EI (1997) *Concise Encyclopedia of Biochemistry and Molecular Biology*, Ed 3. de Gruyter, New York
- Segura S, Scheinvar L, Olalde G, Leblanc O, Filardo S, Muratalla A, Gallegos C, Flores C (2007) Genome sizes and ploidy levels in Mexican cactus pear species *Opuntia* (Tourn.) Mill. series *Streptacanthae* Britton et Rose, *Leucotrichae* DC., *Heliabravoanae* Scheinvar and *Robustae* Britton et Rose. *Genet Resour Crop Evol* 54: 1033–1041
- Smith JAC, Winter K (1996) Taxonomic distribution of Crassulacean acid metabolism. In K Winter, JAC Smith, eds, *Crassulacean Acid Metabolism: Biochemistry, Ecophysiology and Evolution*. Springer, New York, pp 427–436
- Somers DE, Devlin PF, Kay SA (1998) Phytochromes and cryptochromes in the entrainment of the *Arabidopsis* circadian clock. *Science* 282: 1488–1490
- Spiess AN, Feig C, Ritz C (2008) Highly accurate sigmoidal fitting of real-time PCR data by introducing a parameter for asymmetry. *BMC Bioinformatics* 9: 221
- Srivastava GP, Guo J, Shi H, Xu D (2008) PRIMEGENS-v2: genome-wide primer design for analyzing DNA methylation patterns of CpG islands. *Bioinformatics* 24: 1837–1842
- Strayer C, Oyama T, Schultz TE, Raman R, Somers DE, Más P, Panda S, Kreps JA, Kay SA (2000) Cloning of the *Arabidopsis* clock gene TOC1, an autoregulatory response regulator homolog. *Science* 289: 768–771
- Takata N, Saito S, Saito CT, Uemura M (2010) Phylogenetic footprint of the plant clock system in angiosperms: evolutionary processes of Pseudo-Response Regulators. *BMC Evol Biol* 10: 126
- Taybi T, Patil S, Chollet R, Cushnan JC (2000) A minimal serine/threonine protein kinase circadianly regulates phosphoenolpyruvate carboxylase activity in Crassulacean acid metabolism-induced leaves of the common ice plant. *Plant Physiol* 123: 1471–1482
- Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, De Paeppe A, Speleman F (2002) Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biology* 3: RESEARCH0034
- Vidal J, Chollet R (1997) Regulatory phosphorylation of C4 PEP carboxylase. *Trends Plant Sci* 2: 230–237
- Weiss J, Nerd A, Mizrahi Y (1993) Vegetative parthenocarpy in the cactus pear *Opuntia ficus-indica* (L.) Mill. *Ann Bot* 72: 521–526
- Winter K, Garcia M, Holtum JAM (2008) On the nature of facultative and constitutive CAM: environmental and developmental control of CAM expression during early growth of *Clusia*, *Kalanchoë*, and *Opuntia*. *J Exp Bot* 59: 1829–1840
- Wu J, Mao X, Cai T, Luo J, Wei L (2006) KOBAS server: a Web-based platform for automated annotation and pathway identification. *Nucleic Acids Res (Web Server issue)* 34: W720–W724
- Wyka TP, Bohn A, Duarte HM, Kaiser F, Lüttge UE (2004) Perturbations of malate accumulation and the endogenous rhythms of gas exchange in the Crassulacean acid metabolism plant *Kalanchoë daigremontiana*: testing the tonoplast-as-oscillator model. *Planta* 219: 705–713
- Ye J, Fang L, Zheng H, Zhang Y, Chen J, Zhang Z, Wang J, Li S, Li R, Bolund L, et al (2006) WEGO: a Web tool for plotting GO annotations. *Nucleic Acids Res (Web Server issue)* 34: W293–W297